

The Formation of Star Clusters

By BRADLEY C. WHITMORE¹

¹Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD, 21218

The ability of HST to resolve objects ten times smaller than possible from the ground has rejuvenated the study of young star clusters. A recurrent morphological theme found in nearby resolved systems is the observation of young (typically 1 - 10 Myr), massive (10^3 - 10^4 $M_{\odot}$), compact ($\rho \approx 10^5$ $M_{\odot}$ pc⁻³) clusters which have evacuated the gas and dust from a spherical region around themselves. New stars are being triggered into formation along the edges of the envelopes, with pillars (similar to the Eagle Nebula) of molecular gas streaming away from the regions of star formation. The prototype for these objects is 30 Doradus (Figures 1 & 2). Another major theme has been the discovery of large numbers of young (typically 1 - 500 Myr), massive (10^3 - 10^7 $M_{\odot}$), compact star clusters in merging, starbursting, and even some barred and spiral galaxies. The brightest of these clusters have all the attributes expected of protoglobular clusters, hence allowing us to study the formation of globular clusters in the local universe rather than trying to ascertain how they formed ≈ 14 Gyr ago. The prototype is the Antennae Galaxy (Figures 3 & 4).

1. Introduction

1.1. *Hubble's' first six months*

The discovery of spherical aberration in the summer of 1990 raised serious questions about the ability of HST to do the unique science it was designed for, and caused general consternation throughout the astronomical community. Early HST observations of compact star clusters in 30 Doradus and NGC 1275 played pivotal roles in demonstrating that HST, even in its crippled state, could produce stunning results that were impossible to obtain from the ground. This provided a much needed shot in the arm for the Hubble project. In addition, image deconvolution techniques developed to reconstruct the 30 Doradus images showed that most of the compromised resolution could be restored for bright objects.

Before the HST image of 30 Doradus came out in the summer of 1990, several papers had argued that R136 (the central object in the 30 Doradus cluster) was a single star with a mass of several thousand solar masses (e.g., Cassinelli, Mathis & Savage 1981). By the time HST was launched, speckle observations by Weigert & Baier (1985) had resolved the central region into roughly a dozen individual stars. However, concerns about possible artifacts introduced by the speckle technique, and limitations imposed by the small field of view, limited the impact of the results. As the saying goes, “a picture is worth a thousand words”, and it was the spectacular direct images obtained with the WFPC1 on HST that first made it clear just how rich this cluster really was. For example, Hunter *et al.* (1995) identified over 3500 stars in the central $\approx 8''$ alone.

HST was built to make new discoveries, and one of the first was the discovery of young compact clusters in NGC 1275, the central galaxy in the Perseus cluster (Holtzman *et al.* 1992). This demonstrated that HST had no peers when it came to detecting compact, point-like objects against a bright background. Objects that were impossible to see from the ground suddenly became visible. Holtzman *et al.* (1992) also made the rather daring assertion that the young clusters were *protoglobular* clusters formed by a merger event. This was the catalyst for what has become a major cottage industry for Hubble, and provides one of the primary topics for this review.




FIGURE 1. Combined WFPC2 (background) and NICMOS (square inserts) image of 30 Doradus from a 1999 press release by Walborn and Barba. The central star cluster is R136.

1.2. *Motivation*

There are three basic reasons why the HST observations of young star clusters caught the attention of the astronomical community.

1) They provide insight into the mechanisms of star formation, the most fundamental process in astronomy. While we have very detailed models of the structure and evolution of stars (e.g., isocrones in the HR diagram), we have only sketchy ideas of how stars form to begin with. Some of the basic questions that remain to be solved are: Is the initial stellar mass function the same in all environments? Can star formation trigger subsequent star formation? Are all stars formed in groups and clusters?

An obvious approach to solving these questions is to go to where lots of stars are forming, such as merging and starbursting galaxies. When we do this we find that a large fraction of the star formation is in the form of massive, compact star clusters. Understanding how these clusters form should go a long ways toward understanding star formation in general.

2) They allow us to study the formation and evolution of globular star clusters in the local universe rather than trying to ascertain how they formed ≈ 14 billion years ago. An analogy is often made between the study of old globular clusters and the study of

fossils on the earth, since both provide an evolutionary record. Given the opportunity, I believe many paleontologists would switch fields if they could go to where the dinosaurs are still living.

3) They are relevant to the question of whether spiral galaxies can merge to form elliptical galaxies. One of the primary objections to this hypothesis was raised by van den Bergh (1990), who pointed out that spirals have fewer globular clusters per unit luminosity than ellipticals. It now appears that globular clusters can be formed by mergers of gas rich systems (§3.1), providing a natural explanation for this difference.

1.3. *Nomenclature*

For historical reasons, a wide variety of names are currently being used to describe what are physically similar objects. Researchers studying merging galaxies often use names such as “protoglobular cluster” or “young globular clusters”. This is because the focus has been on the question of whether merging galaxies can form globular clusters, in response to van den Bergh’s (1990) criticism of the merger hypothesis for the origin of elliptical galaxies. On the other hand, researchers studying nearby starburst galaxies generally use the term “super star clusters”, a term that was first introduced by Arp & Sandage (1985) when referring to the dominant cluster in the starburst dwarf galaxy NGC 1569. Other names in use include “blue populous clusters” (in the LMC) and “young massive clusters” (in normal spirals; e.g., Larsen & Richtler 1999).

None of these names really ring true, however, which is why one has not become the standard. While there is good evidence that some of the brightest compact clusters become globular clusters, it is obvious that they do not all become globular clusters, since the specific globular cluster frequency in merger remnants would then be too high. The main objection to the term “super star cluster” is that while they may be very luminous for a short period of time, their masses, which are a more fundamental property, are not “super”. They are instead similar to normal globular clusters or open clusters. In essence, there are probably no major physical differences between these various clusters; we are simply seeing young globular clusters, open clusters, and associations at different stages of their evolution or in different environments than we are use to seeing them in the Milky Way.

The defining properties for most of the objects discussed in this review are that they are young, compact, and the brightest are ultra luminous in comparison to old globular clusters. I will simply call them young compact star clusters.

1.4. *Goals*

I have three goals for this review. The first is to highlight the primary contributions that HST has made to the study of the formation of compact star clusters (§2 & 3). There will be an obvious bias towards observations of young compact clusters in merging galaxies, since that is what I know the most about, but I will also highlight some of the major results for nearby clusters such as 30 Doradus. The reader is referred to reviews in this volume by Harris (old globular clusters), Bally (star formation), and Leitherer (starbursts) for related discussions. The second goal is to provide a compilation of the literature on young unresolved compact clusters, as discussed in §4. The third goal is to use the compilation to examine some of the demographics of young cluster formation (§4).

TABLE 1. Properties of Young Compact Star Clusters (approximately in order of distance)

galaxy	N	M_V (bright)	R_{eff}	mass	age	α
near Gal. Center	2	—	—	$10^4 M_\odot$	3 Myr	—
LMC	8	-11.3	2.6 pc	—	3 Myr	—
M82	100	-14.5	3.5 pc	—	100 Myr	—
HE 2-10	76	-12.7	3 pc	$10^3 - 10^5 M_\odot$	—	-1.7
ESO 338-IG04	112	-15.5	—	$10^3 - 10^7 M_\odot$	10-10,000 Myr	—
NGC 1569	7	-13.9	2.2 pc	—	15 Myr	—
NGC 5253	6	-11.1	—	$10^6 M_\odot$	2.5 Myr	—
NGC 1705	36	-13.7	3.4 pc	—	15 Myr	—
NGC 1741	314	-15	—	$10^4 - 10^6 M_\odot$	4 Myr	-1.85
ESO 565-11	700	-13.4	—	—	4 - 6 Myr	-2.2
NGC 1097	88	-13.8	2.5 pc	—	—	—
NGC 4038/39	800	-15.8	4 pc	$10^3 - 10^7 M_\odot$	1 - 500 Myr	-2.1
NGC 3256	1000	-15	5 - 10 pc	—	—	-1.8
NGC 3256 tail	50	-10	—	—	—	—
NGC 3597	700	-13.2	—	—	—	-2.0
NGC 7252	500	-16.2	5 pc	$10^4 - 10^8 M_\odot$	600 Myr	-1.8
NGC 1275	800	-15.8	—	$10^4 - 10^8 M_\odot$	0.1 - 1.0 Gyr	-1.9
NGC 3921	102	-14	< 5 pc	—	500 Myr	-2.1

Notes to Table 1: See Table 2 for references. N is the number of clusters (to $M_V \approx -9$ mag in most cases). M_V (bright) is the M_V magnitude for the brightest cluster. R_{eff} is the average effective radius for the clusters. α is the power-law index for the luminosity function.

2. Nearby Resolved Star Clusters

The techniques for studying star clusters that can be resolved into individual stars are much different than the techniques available for studying more distant clusters. For example, we can determine the stellar luminosity and mass functions, the color-magnitude diagram for the stars, and the radial density profile. For the more distant clusters some of the typical tools are the specific globular cluster frequency, the cluster luminosity function, and color-color diagrams. Before HST, the dividing line between these two regimes was essentially the edge of the Milky Way galaxy. One of the promises of HST was that it would push this dividing line roughly ten times farther away, allowing us to study the clusters in the Magellanic Clouds, and to a lesser extent the nearby galaxies in the Local Group, with the same techniques we have been using for the clusters in the Milky Way. This has indeed been the case. In this section we describe three of the primary HST results for the nearby resolved clusters. Table 1 lists the number, luminosity of the brightest cluster, size, mass, age, and power law index for the luminosity function for a variety of young star clusters that are discussed in this review.

2.1. Young Clusters near the Center of the Milky Way

The Near-infrared Camera and Multi-object Spectrometer (NICMOS) was used by Figer *et al.* (1999) to penetrate the dust toward the center of the Milky Way in order to study two remarkable young clusters near the Galactic Center. Based on turnoffs in the color-magnitude diagrams they estimate that the Arches cluster is 2 ± 1 Myr old and the Quintuplet cluster is 4 ± 1 Myr old. Based on number counts and an extrapolation to 1

$M_{\odot}$, they estimate the masses of the clusters are $\approx 10^4 M_{\odot}$, and the densities are $\approx 10^5 M_{\odot} \text{ pc}^{-3}$.

The existence of these clusters was somewhat unexpected; one would think that the strong tidal shear produced by the central black hole, the high velocity dispersion, and the strong magnetic field would make this a hostile environment for forming young massive clusters. However, given that starburst galaxies are able to support prodigious rates of star formation near their centers, perhaps we should not be surprised that star clusters can also form near the Galactic Center.

Portegies Zwart (2000) performed n-body simulations which indicate that the two clusters should dissolve after 10 - 60 million years in the tidal field of the Galaxy. They also point out that the stellar density near the center of the Galaxy is so high that the clusters would only be distinguishable for a short fraction of their existence, as low as 5 % in some models. Based on this result they conclude that the Galactic Center may be hiding between 10 and 40 clusters which are similar to the Arches and Quintuplet clusters, but slightly older and less compact. Similar simulations by Kim *et al.* (1999) suggest that very massive stars play an important role in the evolution of these clusters because relaxation and mass segregation times are comparable to or even smaller than the lifetimes of the stars.

These clusters are reminiscent of the young clusters found in the central 1.5 kpc of the merger NGC 7252. Miller *et al.* (1997) found ≈ 40 such clusters, all less than ≈ 10 Myr years old based on the (U-B) vs. (V-I) diagram. Hence, it looks like the centers of galaxies may actually be good places to make star clusters, but few if any will survive very long. This suggests that a sizeable fraction of the field star population may have been formed in clusters.

2.2. The Initial Stellar Mass Function (IMF)

The light from young star clusters is dominated by ultraluminous O and B stars. However, if these were the only stars the cluster would not be stable since the massive stars are destined to go supernova, returning their mass to the interstellar medium. Hence, the identification of low mass stars is critical for the survival of the compact star clusters. In addition, the question of whether the IMF is universal, or is instead a function of the environment, is a question which is currently being hotly discussed (see reviews by Scalo 1998, Larson 1999, and Elmegreen 1999). I will briefly comment on a few examples where HST observations of young clusters are relevant to this debate.

The initial stellar mass function is generally parameterized as a power law with an index Γ . The canonical value of Γ for field stars in the Milky Way is -1.35 (Salpeter 1955). There is a lively debate over the question of how universal Γ is, and in particular, whether Γ is the same in clusters and in the field. Several authors have argued that the formation of high mass stars may be favored in starburst regions, which would result in a lower value of Γ .

Initial measurements of the IMF in 30 Doradus were made by Malumuth & Heap (1994; found over 800 stars in the inner $8''$ using the WFPC1), and Hunter *et al.* (1995; found over 3500 stars in the inner $8''$ using WFPC2). More recently, Massey and Hunter (1998) measured the IMF down to $2.8 M_{\odot}$, and find that despite the largest number of high mass and luminosity stars ever seen, the IMF is completely normal. This implies that the IMF is the same over several orders of magnitudes in density, from field stars to starburst clusters like 30 Doradus. The large number of O stars is simply a result of the youth (≈ 2 Myr) and richness of the cluster. A more recent study of WFPC2 images by Sirianni *et al.* (2000) detects stars in 30 Doradus roughly 1 magnitude deeper than Massey and Hunter, and argues that these are pre-main sequence stars in the mass range




FIGURE 2. WFPC2 and NICMOS images in an outlying region of 30 Doradus (i.e., the upper left square inserts in Figure 1) from a 1999 press release by Walborn and Barba. Note the young stars still embedded in dust which are only visible on the NICMOS image (e.g., objects identified as 2, 3, 6 and 7). Also notice the pillar-like dust feature around object 1 pointing away from the central star cluster R136 (see Figure 1 for context).

$0.6 - 3 M_{\odot}$. They construct the IMF in the range $1.35 - 6.5 M_{\odot}$ and find a flattening below $\approx 2 M_{\odot}$. While there are several examples of a flat IMF for stars less massive than $1 M_{\odot}$ (see Larson 1999), this is the first example of a flattening above this point.

Another possible example of a relatively flat IMF, but this time at the high mass end, are the results of Figer *et al.* (1999) for the Arches cluster, one of the two massive young clusters near the Galactic Center. They find $\Gamma = -0.7$ when fitting over the range $6 - 125 M_{\odot}$ for stars in an annulus from $3''$ to $7.5''$. However, incompleteness caused by crowding make this a difficult measurement at the faint end of the mass function. For example, Figer *et al.* note incompleteness fractions of 50 % for stars up to $35 M_{\odot}$ in the inner $3''$. In addition, they find a very flat value of $\Gamma = -0.2$ in the region $2.5''$ to $4.5''$, which may be due to truncation of the faint end by crowding. If we stick to the brighter end of the mass function from 16 to $125 M_{\odot}$, and only use stars in an annulus from $4.5''$ to $7.5''$, $\Gamma \approx -1.1$, still low but not too different from the Salpeter IMF.

Hence, while there is tentative evidence for deviations in the IMF for some environments, the near uniformity over an extremely wide range of environment is quite remarkable. Larson (1999) concludes “this large body of direct evidence does not yet demonstrate convincingly any variability of the IMF, although the uncertainties are still large. Some indirect evidence based on the photometric properties of more distant and exotic systems suggests that the IMF may vary in extreme circumstances, possibly being top-heavy in starbursts and high redshift galaxies.” We also note that low-mass stars are

clearly forming in R136 (e.g., down to $0.6 M_{\odot}$; Sirianni *et al.* 2000), and in other young nearby compact clusters such as NGC 3603 (i.e., no evidence for a flattening down to $1 M_{\odot}$, Eisenhauer *et al.* 1998,). Hence concerns that the young clusters will be unstable due to the absence of low mass stars appear to be unfounded.

2.3. Triggered Star Formation

30 Doradus has been called a “Starburst Rosetta”, since it is the nearest example of a young massive starburst cluster and hence can be studied in unique ways that are not possible for its more distant counterparts. Until recently, 30 Doradus was believed to be a well evolved H II region, with no current star formation going on. Early indications that this was not true were the discovery of an H_2O maser (Whiteoak 1983) and four luminous IR protostars (Hyland *et al.* 1992). More recently, Walborn & Blades (1997) used optical spectral classification of 106 OB stars to show the presence of five distinct stellar populations in 30 Doradus, with ages ranging from ≈ 1 to ≈ 10 Myr. Hence, the simple models of either continuous or single-burst star formation appear to be incorrect. It now appears that a starburst in one area can trigger the formation of stars in a nearby region.

Corroborating evidence for this picture has been obtained by Walborn *et al.* (1999b) using NICMOS, as shown in Figure 1. A roughly spherical shell of “second generation” star formation, including a host of newly discovered IR sources, can be seen around the central R136 concentration (Figure 2). Several massive dust pillars are found streaming away from R136, similar to the famous HST image of the Eagle Nebula. At the heads of these pillars are the sites where active star formation is currently being triggered. The molecular gas revealed by the dust provides the raw material for the star formation. Scowen *et al.* (1998) shows that these pillars are indeed very similar to the pillars in the Eagle Nebula. This same picture of a bright compact central cluster which has evacuated a roughly spherical nebular envelope around it can be seen in a number of the HST press releases (e.g., NGC 604, N11 in the LMC, NGC 4214).

3. Distant Unresolved Star Clusters

In this section we move further out to where it becomes difficult to resolve the individual stars, except in extraordinarily extended clusters such as knot S in NGC 4038/4039 (the “Antennae Galaxies”, Whitmore *et al.* 1999). Historically, most of the early HST observations of compact star clusters were in merging galaxies, followed shortly by similar observations in starburst galaxies. More recently, young compact clusters have also been found in other environments, including barred galaxies, tidal tails, and normal spiral galaxies.

3.1. Young Compact Star Clusters in Merging Galaxies

The primary question for many of the early HST studies of merging galaxies was whether globular clusters were being formed. This possibility was proposed by Schweizer (1987) and Burstein (1987), primarily to address van den Bergh’s (1990) objection to the merger model based on the higher specific frequency of globular clusters in elliptical galaxies. Ashman and Zepf (1992) and Zepf and Ashman (1993) further developed these ideas, and made predictions about the bimodality of the metallicity histogram of globular clusters that should result if most ellipticals are formed by merging spirals.

A hint that young globular clusters might be formed in mergers was provided by Schweizer’s (1982) observations of six unresolved bluish knots in the merger remnant NGC 7252. However, with so few objects he could not be sure they were not simply




FIGURE 3. Image of the Antennae Galaxies (NGC 4038/4039) from Whitmore *et al.* (1999).

field stars. Lutz (1991) observed roughly a dozen blue, point-like objects in the merger remnant NGC 3597, but was not able to resolve the objects and hence could not be certain they were not associations or giant HII regions.

As briefly discussed in §1, the HST observations of NGC 1275 by Holtzman *et al.* (1992) provided the original breakthrough and was the primary catalyst in this field. They discovered a population of about 60 blue compact clusters, and suggested that they were protoglobular clusters which formed ≤ 300 Myr years ago during a merger of NGC 1275 with another galaxy. Unfortunately, NGC 1275, the central cooling-flow galaxy in the Perseus cluster, is such a peculiar galaxy that it is not clear which of its peculiarities is responsible for the formation of the young clusters (e.g., see Richer *et al.* 1993, who suggested that the cooling flows are responsible for the formation of the clusters).

Whitmore *et al.* (1993), using WFPC1 observations of the prototypical merger remnant NGC 7252 (Toomre 1977), found a population of about 40 blue point-like objects with




FIGURE 4. Blowup of two of the brightest clusters in the Antennae (left) and the central regions of the two galaxies (right) from Whitmore *et al.* (1999).

luminosities and colors nearly identical to those found in NGC 1275. Unlike NGC 1275, with all its peculiarities, NGC 7252 is an isolated galaxy which therefore provided a much cleaner connection between the formation of young star clusters and merging galaxies. Whitmore & Schweizer (1995) followed this up with pre-refurbishment observations of another prototypical merger, NGC 4038/4039 (see Figures 3 & 4). Over 700 young star clusters were found in this galaxy. Subsequent observations of both these galaxies using WFPC2 (NGC 7252 - Miller *et al.* 1997; NGC 4038/4039 - Whitmore *et al.* 1999) have increased the numbers of cluster candidates tenfold.

Roughly 30 different gas-rich mergers have now been observed with HST, as summarized in Table 2. In all cases young massive compact clusters have been observed, *the brightest of which have the luminosities, colors, sizes, masses, distributions and spectra that we would expect for globular clusters with ages in the range 1 to 500 Myr.* A few of the key observations are described below, but the reader is referred to the papers listed in Table 2 for the details.

3.1.1. *Luminosities and Colors*

The luminosities of young globular clusters with ages ≈ 10 Myr should be $\approx 5 - 6$ magnitudes brighter than classical old globular clusters, according to the Bruzual and Charlot (1996) models. The models also predict that young globular clusters should be




FIGURE 5. Plot of the evolution in luminosity (ΔV) and in color ($\Delta(V-I)$) of star clusters, based on the Bruzual-Charlot (1996) tracks for a metal-poor population (solid line) and a solar metallicity population (dashed-dot line). The values are normalized to an old, metal-poor population (filled triangle). Ages in Gyr for the solar metallicity track are marked with squares. See the original article for further details (Whitmore *et al.* 1997).

$\approx 1.0 - 1.2$ magnitude bluer in $(V - I)$. Figure 5 shows that this is indeed the case. It also shows how the luminosities and colors of the clusters can be used to age date the clusters. NGC 4038/4039 is clearly the youngest merger remnant, with the mean age of the clusters ≈ 30 Myr and many clusters only a few Myr old. The clusters in NGC 3921 and NGC 7252 are roughly 500 Myr old while NGC 3610 appears to be a 4 ± 2 Gyr merger remnant (Whitmore *et al.* 1997) which may provide the missing link between young mergers and old ellipticals. The case is less certain for NGC 1700, although Brown *et al.* (2000) have recently claimed that this galaxy also has a population of metal-rich clusters that are 3 ± 1 Gyr old. Goudfrooij *et al.* (2000) also found a population of ≈ 3 Gyr clusters in NGC 1316.

3.1.2. Sizes

The ability to measure sizes using HST has been critical to the identification of the young clusters in mergers as candidate globular clusters. Ground based observations, such as those of Lutz (1991), were inconclusive, since they were not able to resolve the clusters to determine whether they were associations or HII regions, with $R_{\text{eff}} \approx 100$ pc, or compact clusters similar to the globular clusters in the Milky Way, with $R_{\text{eff}} \approx 3$ pc (van den Bergh, 1996). Early HST observations using WFPC1 indicated that the

clusters were compact, with $R_{\text{eff}} \approx 10$ pc (Whitmore & Schweizer, 1995). However, van den Bergh (1995) argued that this was too large, and the clusters were more likely to be open clusters. Meurer *et al.* (1995) found that the compact clusters he was studying in very nearby starburst dwarfs were smaller, with $R_{\text{eff}} \approx 2$ pc. He suggested that the apparently larger values in the Antennae were due to poorer resolution and crowding.

Recent observations using the WFPC2 (corrected for spherical aberration) have removed this concern. Several authors have recently measured R_{eff} for young clusters in mergers in the range 3 - 6 pc (i.e., NGC 3921 - Schweizer *et al.* 1996, NGC 7252 - Miller *et al.* 1997, NGC 3610 - Whitmore *et al.* 1997, and NGC 1275 - Carlson *et al.* 1998). Perhaps the best case is for the Antennae galaxies as measured by Whitmore *et al.* (1999), since this is the nearest of the prototypical mergers and the observations were made with subpixel dithering which improves the resolution still further. They find the median effective radii for the clusters is $R_{\text{eff}} = 4 \pm 1$ pc, similar to or slightly larger than those of globular clusters in the Milky Way.

3.1.3. Ages

Ages for the clusters have been estimated in a variety of manners. Figure 5 demonstrates how the luminosities and colors can be used to estimate the ages, as already discussed in §3.1.1. More precise age estimates are possible with more colors, and provide an independent means of solving for the age and the reddening caused by dust. For example, Whitmore *et al.* (1999) use UBVI photometry and reddening-free Q parameters to determine ages for the clusters in the Antennae (Figure 6). They find evidence for four populations of clusters, ranging in age from < 5 Myr to 500 Myr. They also isolate a population of old globular clusters in this galaxy. *Hence, it appears that we can study the entire evolution of globular clusters in this single galaxy.* This is consistent with the simulations of Mihos, Bothun, & Richstone (1993) who find that the merger process takes several hundred million years to complete, hence producing clusters with a wide range of ages.

$\text{H}\alpha$ can be used in two ways to estimate the ages of the younger clusters. The existence of $\text{H}\alpha$ emission itself indicates that a cluster is < 10 Myr, since the O and B stars required to ionize the gas only live for this long (e.g., see the Leitherer & Heckman 1995 models). The second method is to use the size of the $\text{H}\alpha$ ring around a cluster. Whitmore *et al.* estimate that the clusters in the western loop of NGC 4038 are 5 - 10 Myr, since many of them have rings with diameters of $\approx 100 - 500$ pc and measured expansion velocities $\approx 25 - 30 \text{ km s}^{-1}$ (Whitmore *et al.* 1999). The clusters in the overlap region appear to be < 5 Myr old, since the rings are smaller or non-existent in this region.

The most accurate method of estimating ages is to obtain spectra. Zepf *et al.* (1995) obtained spectra of the brightest cluster in NGC 1275 which showed strong Balmer absorption lines, typical of A stars. They estimate ages of 500 Myr for the clusters, although ages from 100 - 900 Myr cannot be ruled out. Schweizer & Seitzer (1998) obtained UV-to-visual spectra of eight cluster candidates in NGC 7252. Six of the clusters have ages in the range 400 - 600 Myr, roughly consistent with the mean photometric age estimate of 650 Myr from Miller *et al.* (1997). One cluster turned out to be an emission-line object with an age estimate of < 10 Myr, indicating that cluster formation is still going on at a low level even in the outer parts of the galaxy. Whitmore *et al.* (1999) obtained GHRS spectra of two clusters in the Antennae with age estimates of 3 ± 1 Myr and 7 ± 1 Myr, in good agreement with the estimates based on the UBVI colors and the $\text{H}\alpha$ morphology.

The youngest clusters appear to be very red objects, which Whitmore & Schweizer (1995) suggested were only now emerging from their dust cocoons. Several of these have




FIGURE 6. Color-color diagram and reddening-free Q parameter diagram for clusters in the Antennae. The numbers on the plots are the values of $\log(\text{age})$. See Whitmore *et al.* (1999) for details.

recently been identified as strong IR sources (Vigroux *et al.* 1996, Mirabel *et al.* 1998, Wilson *et al.* 2000, Gilbert *et al.* 2000, Mengel *et al.* 2000). In fact, the brightest IR source in the Antennae is one of these very red objects (W80), rather than the nucleus of one of the two galaxies. Wilson *et al.* (2000) find three separate molecular clouds around W80 within a region of 1 kpc^2 , and suggest that cloud-cloud collisions may play an important role in cluster formation. However, the lack of similar morphologies for the other very red objects suggest that this may not be the universal mechanism.

3.1.4. Mass

Mass estimates of young compact clusters have been made in two ways. The first is based on the luminosity and color of the clusters using stellar population models such as Bruzual & Charlot (1996). These estimates generally range from 10^3 to $10^7 M_{\odot}$ (see Tables 1 and 2), in good agreement with old globular clusters with a mean of $2 \times 10^5 M_{\odot}$ (Mandushev, Spassiva & Staneva (1991). A more direct method of determining the mass is to measure the velocity dispersion of the stars in the clusters. Observations have

been obtained for nine clusters so far (two in NGC 1705 and one in NGC 1569 by Ho and Fillipenko 1996; two in M82 by Smith and Gallagher 2000, 4 in NGC 4038/39 by Mengel *et al.* 2000). The dispersions range from 10 - 20 km s⁻¹ and the masses range from 1×10^5 to 4×10^6 M_⊙, in good agreement with values for the more massive old globular clusters in the Milky Way. The size and dispersion measurements also show that the crossing times for the clusters are ≈1 Myr. Hence, even the younger clusters have survived many crossing times. The older clusters in NGC 7252, NGC 3921, NGC 4038/4039, and NGC 1275 (≈ 500 Myr; see Tables 1 and 2), have survived for several hundred crossing times and appear to be quite stable. Their densities are ≈ 10⁵ M_⊙ pc⁻³, similar to old globular clusters, hence these clusters will almost certainly last for tens of Gyr.

3.1.5. *The Luminosity Function*

To first order, the luminosity functions of young compact clusters in merging galaxies are power laws with index ≈ -2 (Table 1). Harris and Pudritz (1994) have pointed out that the mass function for giant molecular clouds is also a power law with a similar index. Hence, all that may be necessary is a triggering mechanism to get the molecular clouds to collapse and form star clusters. Jog & Solomon (1992) have suggested that merger induced starbursts can raise the ambient pressure in the ISM and trigger the implosion of the molecular clouds. Elmegreen & Efremov (1997) agree that high-pressure environments are needed to trigger the star formation and suggest other mechanisms might be high background virial density (e.g., in dwarf galaxies), turbulent compression, or large-scale shocks (in interacting galaxies).

The power law index for the young clusters is markedly different than the Gaussian profile found for old globular clusters (e.g., Figure 3 of Zhang and Fall 1999). However, various destruction mechanisms (e.g., 2-body evaporation, bulge and disk shocking, dynamical friction, stellar mass loss) should modify the distribution with time. Two-body evaporation appears to be the strongest amongst these mechanisms, destroying the fainter more diffuse clusters first, and in certain conditions leaving a peaked distribution similar to what is seen for old globular clusters (e.g., Fall and Zhang, 2000). This is similar to young clusters in the Milky Way with the OB associations typically only lasting tens of Myr, and open clusters lasting hundreds of Myr. Other examples of clusters which are apparently dissolving are the Arches and Quintuplet clusters near the Galactic Center, since no older clusters are seen in their vicinity, and the ≈ 40 clusters in the inner 6'' of NGC 7252, which all have ages less than about 10 Myr (Miller *et al.* 1997). Finally, the number of young clusters in the Antennae galaxies is so large that it requires most of the clusters to dissolve or the value of S_N will be too high when it settles down to become an elliptical (Whitmore *et al.* 1999)!

Fritze-v.Alvensleben (1999), following a similar line of reasoning to Meurer (1995), has attempted to determine the mass function for the clusters in the Antennae using the color information from the WFPC1 observations by Whitmore & Schweizer (1995). They conclude that the original mass function is a Gaussian which gets spread out in time to form the power law luminosity function we observe today. However, their analysis does not take into account the fact that the cutoff in the observed luminosity function is due to incompleteness at the faint end (see Zhang & Fall 1999 for a discussion). When convolved with uncertain age estimate used to convert from luminosity to mass (e.g., due to reddening from dust and the availability of only V-I colors), the resulting distribution will artificially appear to be roughly Gaussian. A more complete treatment by Zhang & Fall (1999), using UBVI colors based on WFPC2 observations by Whitmore *et al.*




FIGURE 7. Luminosity of cluster candidates in the Antennae as a function of their ages (from Zhang and Fall, 1999, Fig. 2). The lines represent the Bruzual-Charlot (1996) tracks with $\log(M/M_\odot) = 6.0$ (top), 5.5, 5.0, 4.5, and 4.0. See Zhang and Fall for further details.

(1999) and corrections for reddening and incompleteness, concludes that the mass function is roughly a power law.

There is some evidence that the luminosity function for the young clusters is not a perfect power law, but is steeper for bright magnitudes (NGC 4038/4039 - Whitmore *et al.* 1999, NGC 3256 - Zepf *et al.* 1999). In the Antennae, Whitmore *et al.* (1999) find that the cluster luminosity function appears to have a bend at $M_V \approx -10.4$ (≈ -11.4 after making a correction for extinction). For absolute magnitudes brighter than $M_V \approx -10.4$ the power law is steep and has an exponent of $\alpha = -2.6 \pm 0.2$, while for the range $-10.4 < M_V < -8.0$ the power law is flatter, with $\alpha = -1.7 \pm 0.2$. Assuming a typical age of 10 Myr for the clusters, and 1 mag of extinction, the apparent bend in the LF corresponds to a mass $\approx 1 \times 10^5 M_\odot$, only slightly lower than the characteristic mass of globular clusters in the Milky Way. A similar bend may be present in the mass function derived by Zhang & Fall (1999). The bend may be a precursor to what will become the peak of the globular cluster luminosity function.

3.2. Young Compact Star Clusters in Starburst Galaxies

Young compact star clusters are also found in many starburst galaxies, but in much smaller numbers than the merging galaxies. Meurer *et al.* (1992), using ground-based observations, found a population of young clusters in the nearby starburst dwarf galaxy NGC 1705, the brightest of which was an unresolved off-center nucleus which they proposed as a young (13 Myr) globular-like cluster with a mass $\approx 1.5 \times 10^6 M_\odot$. Meurer *et al.* (1995) followed this up with an extensive study of nine starburst galaxies obtained

with the Faint Object Camera on HST. All nine of the galaxies contained young compact star clusters. On average, 20 % of the UV light from the galaxies comes from the clusters. The brightest clusters are preferentially found near the centers of the galaxies. They find the sizes are similar to Galactic globular clusters and the luminosity function has an index ≈ -2 . Hence, the clusters found in the starburst galaxies appear to be similar to the clusters found in merging galaxies.

Several other authors find similar examples in other starburst galaxies, as listed in Table 3. Conti & Vacca (1994) observed 19 “knots” in the Wolf-Rayet galaxy He 2-10, each with a luminosity, mass, and size similar to Galactic globular clusters. Other early observations include those of O’Connell *et al.* (1994) for NGC 1569 and NGC 1705, Hunter *et al.* (1994) for NGC 1140, and Watson *et al.* (1996) for NGC 253.

The case of M82, the prototypical starburst dwarf galaxy, is especially interesting. O’Connell *et al.* (1995) find a complex of over 100 compact, luminous “super star clusters” concentrated in the inner 100 pc of the galaxy shining through a relatively dust free region. The brightest cluster has $M_V = -13.2$ while the mean M_V is -11.6 mag. Since most of this galaxy is embedded in dust the total number of young clusters is likely to be several times this value. It is quite possible that the starburst in M 82 was triggered by a tidal interaction with its larger neighbor, M 81, hence it is not clear whether M 82 (or several other starburst galaxies with evidence for interactions) should be in this section or in the previous section on interacting galaxies. De Gris, O’Connell, & Gallagher (1999) studied a region farther from the center of M82 where active star formation is not occurring. They estimate the ages of the clusters in this region at 20 - 100 Myr. They find that the objects in the outer regions have sizes in the range $2.3 < R_{\text{eff}} < 8.4$ pc. While the lower value is similar to the sizes of galactic globular clusters, the higher value is more typical of open clusters. They also find that the brightest clusters have $M_V \approx -10$ mag, and most are in the range -5 to -7 mag. Hence, most of these clusters are too faint to become globular clusters since they will fade several magnitudes as the stars evolve. It appears that this region is not able to form the true “super star clusters” seen near the center of M82 and in other merger and starburst systems.

The lesson appears to be that luminous young star clusters are found whenever there is vigorous star formation, whether it be in mergers or starburst galaxies. Since the ultraluminous IRAS sources are essentially all mergers (Sanders *et al.* 1988), it is not surprising that mergers show the largest populations of young star clusters.

3.2.1. *Young Compact Star Clusters in Barred Galaxies*

Barth *et al.* (1995) found young clusters in the circumnuclear star-forming rings around the barred spiral galaxies NGC 1097 and NGC 6951. The clusters are compact, with $R_{\text{eff}} \approx 2.5$ pc in NGC 1097 and ≤ 4 pc in NGC 6951. The brightest cluster has $M_V(\text{uncorrected for extinction}) = -12.6$. They estimate an intrinsic M_V in the range -14 to -15 , since the clusters are on the outer edges of prominent dust lanes. Hence, these clusters appear to be quite similar to clusters in merging and starbursting galaxies. Maoz *et al.* (1996) found a similar population of clusters in a sample of five barred galaxies using the Faint Object Camera. They estimate that 10 - 40 % of the UV light is from compact clusters in these galaxies, similar to the result for starburst galaxies (Meurer *et al.* 1995). Buta *et al.* (2000) have done a careful analysis of young compact clusters in the nuclear ring of NGC 1326, an early-type barred spiral in the Fornax cluster. They find 269 candidate clusters with ages in the range 5 to 200 Myr, but no clusters older than this. The older clusters still lie within the ring, with no evidence of migration. The luminosity function has an index of -2.1 , similar to the other compact clusters discussed in this review, but the brightest clusters are fainter than the brightest clusters found in

mergers or starburst galaxies, with no M_V (uncorrected) brighter than -11. The authors conclude that this galaxy lacks any true super star clusters, and suggest that super star clusters are not a universal property of star-forming rings. It is interesting to note that while the strong Lindblad resonance in this galaxy can produce clusters with a typical power-law luminosity function, it apparently cannot form the brightest clusters which are the best candidates for protoglobular clusters.

3.3. Young Star Clusters in Spiral Galaxies

Larsen & Richtler (1999) carried out a systematic ground-based search for young massive clusters in 21 nearby non-interacting spiral galaxies and found young massive star clusters in about one quarter of the galaxies. In a followup paper (Larsen & Richtler 2000), they add a variety of other galaxies to the sample from the literature, including merging and starbursting galaxies, in order to test what conditions are most advantageous for making large numbers of massive clusters. They define the *specific cluster frequency* (not to be confused with the specific globular cluster frequency, S_N , see Harris 1991) as the fraction of light in clusters to the fraction of light in the total galaxy: $T_L = 100 \times L_{clusters} / L_{galaxy}$.

They prefer to make the measurement in U which is most sensitive to young clusters. Their primary result is that $T_L(U)$ is well correlated with the star formation rate per unit area (Figure 8). Galaxies with very active star formation form proportionally more of their stars in clusters than in the field, with some merger and starburst galaxies devoting as much as 15 - 20 % of their luminosity to clusters. Note that this is precisely what is needed to increase the specific globular cluster frequency, a concern voiced by Harris (1999). Larsen & Richtler (2000) also argue that “The cluster formation efficiency seems to depend on the SFR in a continuous way, rather than being related to any particularly violent mode of star formation”.

Closer to home, Chandar *et al.* (1999) have used WFPC2 observations to study the young compact clusters in M33. They find 44 young clusters with ages ≤ 100 Myr and masses in the range 6×10^2 to $2 \times 10^4 M_\odot$. Hence, M33 appears to currently be making many young compact clusters, but few if any with the masses of regular globular clusters.

3.3.1. Young Compact Star Clusters in Tidal Tails

Knierman *et al.* (2000) have examined six tidal tails in four prototypical mergers (NGC 3256, NGC 3921, NGC 4038/4039 and NGC 7252). They find that only one of the tails (the western tail of NGC 3256) currently has a large number of young compact clusters (i.e. ≈ 50 clusters with the brightest having $M_V \approx -10$ mag). It is not clear whether the clusters were formed when the tidal tail was ripped from the galaxy or are currently forming. Some of the other tails appear to have only a few young clusters, (e.g., NGC 7252 and NGC 3921) while others (e.g., NGC 4038/39) appear to have essentially no clusters in the tails. Hence, it appears that there is a wide range in the number of clusters in tidal tails, perhaps due to differences in how the tails were generated (e.g., gas-rich versus gas poor, deep penetrating orbit versus quiescently being pulled out from the outer regions of the galaxy, etc.). Other studies including observations of young compact star clusters in tidal tails include Lee, Kim & Geisler (1997), Tyson *et al.* (1998), and Gallagher *et al.* (2000).




FIGURE 8. Plot of the specific luminosity for the clusters in the U band vs. the star formation rate per unit area for a sample of spiral, starburst, and merging galaxies (from Larsen & Richtler *et al.* 2000, Fig 6).

4. A Compilation of the Literature and a Discussion of Broader Issues

Tables 2, 3, and 4 provide a compilation of HST (and occasionally key ground-based) observations of young compact star clusters in merging, starbursting, and miscellaneous other galaxies.

Based on the discussion in §2 and 3, and the compilation in Tables 1, 2, 3, and 4, it is clear that luminous young compact star clusters are produced in a wide variety of environments, but in much greater number in mergers and starburst galaxies, systems where vigorous star formation is occurring. A similar conclusion was reached by van den Bergh (2000), who comments, “Presently available data strongly suggest that the specific cluster forming frequency is highest during violent bursts of star formation”. In addition, the most luminous clusters are formed in the regions with the most violent star formation.

An important question is whether this is a statistical effect due to the lower number of clusters in galaxies with low star formation, or whether it is physically more difficult to form massive clusters in relatively quiescent systems (i.e., “is there a cutoff at the high end of the luminosity function for quiescent galaxies?”). The situation may be analogous to the upper IMF in 30 Doradus. It was presumed that the large number of very luminous stars indicated that conditions in 30 Doradus were especially conducive for making high mass stars. However, Massey & Hunter (1998) find that the IMF is normal; that the large number of massive stars is simply due to the tremendous number of stars in the system and the young age of the cluster.

The most straightforward approach to answering this question would be to look at the mass function of the clusters for a variety of galaxies. Unfortunately, this is quite difficult given the large amounts of dust and the dimming caused by stellar evolution. The only galaxy where this has been attempted in detail is the Antennae (Zhang & Fall




FIGURE 9. Approximate luminosity functions for galaxies in Table 1, normalized to have 0.25 mag bins.

1999). However, we can attempt to make the comparison using the luminosity function, as shown in Figure 9 for 8 galaxies. We find that all the galaxies have luminosity functions with similar slopes, with an average power law index $\alpha = -1.93 \pm 0.06$ (uncertainty in the mean; the scatter is 0.18). The primary difference is the normalization of the luminosity function, with NGC 3256 and NGC 4038/39 having large numbers of clusters while NGC 3921 and HE 2-10 have relatively few clusters. There is no obvious trend for a cutoff at high luminosity for the more quiescent galaxies, suggesting a universal luminosity function is a reasonable approximation.

Such an approach is oversimplified for a number of reasons, primary amongst them being that the luminosity of the clusters vary with time. For example, a single-age burst population will evolve to the right in Figure 9, making it difficult to determine whether the luminosity function is lower because of evolution or due to a smaller number of clusters originally forming. Other difficulties with this simple model are that it assumes similar star formation histories for the various galaxies (e.g., continuous rather than sporadic bursts at different times), and ignores the fact that the faint end will probably undergo rapid evolution as the faint clusters dissolve. Nevertheless, to first order the luminosity functions appear to be remarkably similar in form.

Another approach which allows us to increase the sample at the expense of more scatter for any particular galaxy is to plot the magnitude of the brightest cluster vs. the number of clusters in the galaxy, as shown in Figure 10. This figure uses the groundbreaking survey of Larsen & Richtler (2000), with the additions of some new points for merging

TABLE 2. Observations of Interacting Galaxies with Young Star Clusters

Reference	Brief Description
Schweizer (1982)	NGC 7252 (ground-based, 6 knots, stat. significant?)
Lutz (1991)	NGC 3597 (ground-based, ≈ 10 knots, lacked resolution)
Holtzman et al. (1992)	NGC 1275 (proposed “protoglobular clusters”, $n = 60$)
Whitmore et al. (1993)	NGC 7252 (prototypical merger, $n = 40$)
Crabtree et al. (1994)	NGC 7727 (ground-based)
Whitmore & Schweizer (1995)	NGC 4038/4039 ($n = 700$, Antennae galaxies)
Zepf et al. (1995)	NGC 1275 (ground-based spectra, .1 - 1 Gyr)
Borne et al. (1996)	Cartwheel galaxy (clusters in rings)
Holtzman et al. (1996)	NGC 3597, NGC 6052 (mergers, not cooling flows)
Schweizer et al. (1996)	NGC 3921 (102 candidate globulars, 49 “associations”)
Hilker & Kissler-Patig (1996)	NGC 5018 (several hundred Myr to 6 Gyr)
Miller et al. (1997)	NGC 7252 ($n = 499$, 3 pop., < 10 Myr for $R < 6''$)
Whitmore et al. (1997)	NGC 1700, NGC 3610 (missing link with ellipticals ?)
Schweizer & Seitzer (1998)	NGC 7252 (spectra, $n = 8$, ages, metallicities)
Brodie et al. (1998)	NGC 1275 (ground-based spectra, age ≈ 450 Myr)
Carlson et al. (1998)	NGC 1275 ($n = 3000$, mix of red and blue clusters)
Johnson et al. (1998)	NGC 1741 (starburst, interacting, Hickson group)
Stiavelli et al. (1998)	NGC 454 (5-10 Myr, effects of emission on photometry)
Dinshaw et al. (1999)	NGC 6090 ($n = 4$, NICMOS observations)
Zepf et al. (1999)	NGC 3256 ($n = 1000$, 15-20 % of U light, break in LF ?)
Whitmore et al. (1999)	NGC 4038/4039 ($n=800$ to 8000, break in LF ?)
Gallagher et al. (2000a)	Stephan’s Quintet ($n = 150$, galaxies and tidal tails)
Alonso-Herrero et al. (2000)	Arp 299 (ULIRG, $n = 40$)
Forbes & Hau (2000)	NGC 3597 (ground-based, K band, $\alpha = -2$)
Johnson & Conti (2000)	HCG 31 (several in Hickson Compact Group 31)
Gilbert et al. (2000)	NGC 4038/39 (IR spectra, ages, masses)
Mengel et al. (2000)	NGC 4038/39 (IR spectra, ages, masses)
Georgakakis et al. (2000)	NGC 6702 (dust-lane elliptical, 2- 5 Gyr)
Goudfrooij et al. (2000)	NGC 1316 (elliptical with shells, 3 Gyr)

and starburst galaxies drawn from the papers in Tables 2, 3, and 4. We find a clear trend between the number of clusters observed and the magnitude of the brightest cluster. The solid line is the fit to the data (excluding NGC 1569) with a slope = -2.3 ± 0.2 . The dotted line shows the trend expected if there is a universal luminosity function with $\alpha = -2$ and the increase in the luminosity is simply due to a larger sample of clusters (i.e., the slope is -2.5). Again, to first order it appears that a universal luminosity function can explain the data, even with the large scatter expected from low number statistics, non-uniform databases, differences in selection criteria, and differences in cutoff magnitudes (only those with cutoffs ≈ -9 have been included).

However, this may not be the whole picture. It is easy to think of examples that do not appear to fit this picture. For example, the brightest clusters in NGC 1569 and NGC 1705 are 2 - 3 magnitudes brighter than the second brightest cluster in the galaxy (O’Connell, Gallagher & Hunter 1994, see Figure 9 of Meurer *et al.* 1995 which provides a graphical representation for NGC 1705), suggesting something special is happening in these clusters. In addition, many galaxies are currently forming a few very young compact clusters (e.g, the central regions of the Milky Way and NGC 7252) which, assuming a steady formation rate over a long period of time, implies that a few very massive clusters

TABLE 3. Observations of Starburst Galaxies with Young Star Clusters

Reference	Brief Description
Arp & Sandage (1985)	NGC 1569 (ground-based, coined “super star clusters”)
Kennicutt & Chu (1988)	LMC (cores of HII regions may be globular clusters)
Meurer et al. (1992)	NGC 1705 (ground-based, $10^6 M_{\odot}$)
Conti & Vaca (1994)	He 2-10 (Wolf-Rayet galaxy, 1 - 10 Myr, $10^5 - 10^6 M_{\odot}$)
Hunter et al. (1994)	NGC 1140 ($n = 7$, merger ?, 3 - 15 Myr)
O’Connell et al. (1994)	NGC 1569, 1705 ($n=3$, $R_{\text{eff}} \approx 3$ pc, density $\gg R136$)
Meurer et al. (1995)	9 starbursts (20 % of UV from clusters, $\alpha = -2$)
O’Connell et al. (1995)	M82 ($n \approx 100$, $R_{\text{eff}} = 3.5$ pc, near center)
Watson et al. (1996)	NGC 253 ($n = 4$, brightest = -15 mag and $1.5 \times 10^6 M_{\odot}$)
Leitherer et al. (1996)	NGC 4214 (FOC and FOS, $n = 200$, 4-5 Myr)
de Marchi et al. (1997)	NGC 1569 (1569A is superposition of two clusters)
Ho & Fillppenken (1997)	NGC 1705, NGC 1569 (spectra, velocity disp., $3.3 \times 10^5 M_{\odot}$)
Calzetti et al. (1997)	NGC 5253 (BCG, $n = 6$, 2.5 Myr, $\approx 10^6 M_{\odot}$)
Oslin et al. (1998)	ESO338-IG04 (BCG)
De Grijs et al. (1999)	M82 (outer region, 20 - 100 Myr, fainter than -10 mag)
Gallagher et al. (2000)	NGC 7673, NGC 3310, Haro I (clumps of SSCs)
Johnson et al. (2000)	He 2-10 (Wolf-Rayet galaxy, WFPC2, H_{α} , GHRS)
Smith et al. (2000)	M82 (ground-based spectra, 60 Myr, $2 \times 10^6 M_{\odot}$)
Oslin (2000)	Mrk 930, ESO185-IG13, ESO350-IG38 (BCGs)
Meurer (2000)	NGC 3310 (0 - few 100 Myr, continuous formation)

TABLE 4. Observations of Other Galaxies with Young Star Clusters

Reference	Brief Description
Barth et al. (1995)	NGC 1097, NGC 6951 (barred, 2 - 3 pc, $n = 88$ and 24)
Holtzman (1996)	Abell 496, 1795, 2029, 2597 (not related to cooling flows)
Maoz et al. (1996)	5 barred (FOC observations, 10 - 40 % of UV from clusters)
Lee et al. (1997)	UGC 7636 (dwarf near NGC 4472, $n = 18$, tidal tail)
Carollo et al. (1997)	35 spirals (M_V vs. R_e diagram)
Tyson et al. (1998)	NGC 5548 (in tidal tail of Seyfert galaxy)
Buta et al. (1999)	ESO 565-11 (barred, $n = 700$, 4-6 Myr, $\alpha = -2.2$)
Chandar et al. (1999)	M33 ($n=44$, $10^3 - 10^4 M_{\odot}$, < 100 Myr)
Buta et al. (2000)	NGC 1326 (barred, $n = 269$, 5 - 200 Myr)
Larsen & Richtler (2000)	21 spirals (ground-based spirals, also mergers & starbursts)
Gallagher et al. (2000)	Stephans’ Quintet (Hickson compact group)
Knierman et al. (2000)	Tidal tails in 4 mergers (not all have clusters)

should eventually form, based on the statistics. These appear to be missing. However, it is possible that the young clusters are forming in regions that are not conducive to the long-term survival of the clusters.




FIGURE 10. Plot of the magnitude of the brightest cluster vs. the log of the number of clusters. Filled circles are spiral galaxies from Larsen & Richtler (2000), open circles are mergers, stars are starbursting galaxies, and the half filled square is a barred galaxy (Table 1). The solid line is a best fit (excluding NGC 1569) while the dashed line is the prediction from a universal power law luminosity function with index $\alpha = 2$. See text for more details.

5. Current and Future Questions

The HST observations of star clusters have answered many questions, but typical of any active field of science, they have introduced even more new questions. Here are some of the fundamental questions that should be addressed over the coming decade.

5.1. *Will some of the young compact clusters survive to become classical globular clusters, and if so, how many ?*

Historically, astronomers have been approaching this question from two different directions. Looking at resolved clusters, Kennicutt and Chu (1988) concluded “that populous clusters may be forming in giant HII regions, but only a small fraction of giant HII regions are likely to contain such clusters”. The prototype for this idea is 30 Doradus. For the more distant galaxies where the clusters are not well resolved, most of the original motivation came from trying to understand the specific frequency of globular clusters in elliptical galaxies (Schweizer 1982, Burstein, 1982, Ashman and Zepf 1992). As pointed out in §3.1.5, we actually need the vast majority of clusters to dissolve or we end up with specific globular cluster frequencies that are too high.

It now seems well established that some of the young compact clusters will survive to form globular clusters. For example, in NGC 7252 and NGC 3921 the clusters are

already 500 Myr (several hundred crossing times), have the distribution expected of globular clusters, and have the same masses and densities of classical globular clusters. The remaining question is *how many* of the clusters will survive and become old globular clusters. In particular, is this how the red (metal-rich) population of globular clusters found in elliptical galaxies are formed, as Ashman & Zepf (1992) propose, or is this just a minor trace population? For example, Schweizer *et al.* (1996) concludes that the total number of globular clusters in NGC 3921 has increased by only 40 %. While sizeable, this may not be enough to explain the increase in S_N in ellipticals unless the typical ellipticals has several major mergers in its lifetime. In addition, a clear prediction from Ashman & Zepf is that the ratio of red (metal-rich) to blue clusters should increase for high S_N galaxies. However, Forbes, Brodie, & Grillmair (1997) find that the number of red clusters does not increase with S_N . It is possible that this is because their sample is dominated by ellipticals in clusters of galaxies, where other mechanisms might also be operating (e.g., stripping the globular clusters out of nearby dwarf galaxies; see the Harris review in this volume). In summary, it appears that many of the brighter young compact clusters will become classical globular clusters, but the jury is still out on whether this is the cause of the increase in S_N for elliptical galaxies.

5.2. *What fraction of stars are formed in clusters?*

Since only a subsample of the young clusters are likely to survive, an obvious question is whether most of the field stars in a galaxy are originally formed in clusters. In the Milky Way, approximately 0.1% of the stars are currently in globular clusters. However, in some starbursting and merging galaxies the fraction of light from the clusters is as high as 20 % (Meurer *et al.* 1995). Even in these young star forming regions many of the field stars are from clusters that have already dissolved, hence the true percentage of stars that were originally in clusters is even higher, and might conceivably be ≈ 100 %.

HST observations may allow us to answer this question by determining the rate at which clusters dissolve. For example, if we were to assume that the Antennae has been making clusters at the same rate for the past 200 Myr (a rather uncertain assumption to say the least), we could use figure 2 from Zhang and Fall (reproduced as Figure 7) to show that for every 20 clusters originally formed, only about 1 will survive to an age of ≈ 100 Myr (i.e. there are roughly the same number of clusters in the 0 - 10 Myr age bin as in the 20 - 200 Myr age bin). While this very crude calculation is probably not justifiable for a single galaxy, which we may be catching during the peak of cluster formation, once a larger sample becomes available this would be a reasonable approach.

An intriguing result is the finding that the luminosity function in the Ursa Major dwarf spheroidal galaxy and the globular cluster M15 are essentially indistinguishable (Wyse *et al.* 1999), even though the densities differ by three order of magnitude. It is tempting to suggest that perhaps most the stars are formed in groups and clusters, and that the field stars are simply the remnants of the fainter, less dense clusters which have dissolved.

5.3. *What fraction of star formation is triggered by other star formation?*

There appears to be a variety of ways to form stars (e.g., gravitational instabilities, shocks between colliding clouds of gas, enhanced pressure of the ISM, etc.). As discussed in §2.3, HST observations suggest that star formation can also be triggered by nearby bursts of star formation (e.g., around 30 Doradus; see Figures 1 and 2). An interesting question is what fraction of all stars have been formed this way? At any one time only a relatively small fraction of star formation appears to be triggered (e.g., the clusters around 30 Doradus are relatively modest compared to 30 Doradus itself). However, in principle this is a self-propagating process which may continue over a much longer period

of time, hence it is possible that overall a relatively large fraction of star formation is triggered. The fact that much of the triggered star formation is still embedded in dust clouds makes it difficult to obtain a complete census. New observations with the NICMOS + cryocooler, and the IR channel of the WFC3, will help answer the question of how important this mechanism is to the total production of stars in a galaxy.

5.4. *Is a massive open cluster the same as a low-mass globular cluster ?*

The fact that the luminosity functions for young clusters are power laws begs the question of whether there is anything fundamentally different between a massive open cluster and a low-mass globular cluster. Are we looking at a continuum, or a bimodal distribution with fundamentally different formation mechanism for open clusters and globular clusters? It seems possible that the distinction between the two is artificial, and is due to the fact that we live in a galaxy which had an initial burst of star formation 14 Gyr ago but no major bursts since then. The only clusters that have survive from the initial burst are, by necessity, massive and compact. These we call globular clusters. In the present epoch, the star formation rate is percolating at a much lower rate and we are only able to see the spectrum of clusters from associations to open cluster. We do not see young globular clusters for several possible reasons. First, they should only form very rarely, since the star formation rate is so low (see §4). Second, we would probably call them open clusters anyway, since we are not use to calling anything young a globular cluster. Indeed, there is overlap in the masses of open and globular clusters. Candidate open cluster/globular clusters might include M67 (5 Gyr), Be 17 and Lynga 7, which according to Phelps *et al.* (1994), may be as old as the youngest globular clusters.

5.5. *Can we develop a unified picture of cluster formation that explains all this?*

While we are making good progress understanding many pieces of the puzzle, how it all fits together is a much tougher question. Is it possible to develop a universal model that provides a framework for understanding cluster formation both near (e.g., the classic picture of associations, open clusters, and globular clusters in the Milky Way), intermediate (“populous” star clusters in the LMC and “super star clusters” in nearby dwarf starbursts), and far (the young compact clusters in mergers and starbursts); for spiral and elliptical galaxies; for the initial collapse of a many galaxies ≈ 14 Gyr ago and mergers of galaxies in the local universe; and for violent starbursts and “quiescent” star formation? Some of the ideas discussed in this review lead to the following working hypothesis, portions of which several groups are pursuing, in particular Elmegreen & Efremov (1997), Vesperini (1998), and Fall & Zhang (2000).

The luminosity functions for young clusters (e.g., Whitmore & Schweizer 1995), molecular clouds (Harris & Pudritz 1994) and HII regions (Elmegreen & Efremov 1997) are all power laws with index ≈ -2 . Hence, we start with a universal power law luminosity (mass) function for. The total number of clusters is normalized depending on how active the star formation is (e.g., Larsen & Richtler, §4). This explains the existence of large numbers of bright clusters in mergers, since they have the most active star formation. The physics of how the cluster formation is triggered is still uncertain, but several possible mechanisms have been proposed (e.g., Jog & Solomon 1992, Kumai *et al.* 1993, Elmegreen & Efremov 1997). The power law evolves due to both internal (e.g., evaporation, stellar mass loss) and external (e.g., tidal stress, triggered star formation) influences (e.g., Fall & Zhang 2000), with most of the faint and/or diffuse clusters dissolving, just as they do in the Milky way. This model would then need to be convolved with models of galactic evolution (i.e., a combination of initial collapse, hierarchial merging, and in-

ternal galactic dynamics), and stellar evolution (dimming and reddening of the starlight) to produce the wide variety of cluster demographics we see in galaxies.

5.6. *What is the limiting redshift for observing young globular clusters with HST and NGST?*

The current limiting redshift for observing young globular clusters with characteristics similar to the brightest young clusters in the Antennae is $Z \approx 0.5$, using the WFPC2 on HST. It will be possible to do slightly better (i.e. $Z \approx 0.8$), with the Advanced Camera for Surveys since it has a quantum efficiency which is roughly 3 times better than WFPC2. However, the real breakthrough will be NGST, where Burgarella & Chapelon (1998) estimate that it will be possible to observe young globular clusters out to $Z \approx 9$, if they exist. Since globular clusters appear to be the oldest fossils we observe in galaxies it is quite possible that the first objects we will see emerging from the “dark ages” will be young compact star clusters, similar to what we are seeing in nearby galaxies.

REFERENCES

- ARP, H. & SANDAGE, A. 1985 *AJ* **90**, 1163.
- ASHMAN, K. M. & ZEPF, S. E. 1992 *ApJ* **384**, 50.
- ALONSO-HERERO, A., RIEKE, G. H., RIEKE, M. J., & SCOVILLE, N. Z. 2000, *astroph/9911534*.
- BARTH, A. J., HO, L. C., FILIPPENKO, A. V., & SARGENT, W.L.W. 1995 *AJ* **110**, 1009.
- BORNE, K. D. ET AL. 1996 *Science with the Hubble Space Telescope - II*, eds. P. Benvenuti, F. D. Macchetto, & E. J. Schreier, 239.
- BRODIE, J. P., SCHRODER, L. L., HUCHRA, J. P., PHILLIPS, A. C., KISSLER-PATIG, M., & FORBES, D. F. 1998 *AJ* **116**, 691.
- BROWN, R. J. N., FORBES, D. A., KISSLER-PATIG, M., & BRODIE, J. P. 2000 *MNRAS* in press.
- BRUZUAL A. G., & CHARLOT, S. 1996, unpublished.
- BURGARELLA, D. & CHAPELON, S. 1998 *Science with the NGST*, ed. E. P. Smith & A. Koratkar (ASP Series Vol. 133), 227.
- BURSTEIN, D. 1987 *in Nearly Normal Galaxies*, ed. S. Faber (Springer, New York) **47**.
- BUTA, R., CROCKER, D. A. & BYRD, G.G. 1999 *AJ* **118**, 2071.
- BUTA, R., TREUTHARDT, P. M., BYRD, G.G. BYRD, & CROCKER, D. A. 2000 *AJ*, in press.
- CALZETTI, D., MEURER, G. R., BOHLIN, R. C., GARNETT, D. R., KINNEY, A. L., LEITHERER, C., STORCHI-BERGMANN, T. 1997, *AJ*, **114**, 1834.
- CARLSON, M. N. & THE WFPC2 TEAM 1998 *AJ* **115**, 1778.
- CAROLLO, C. M., STIAVELLI, M., DE ZEEUW, P. T. & MACK, J. 1997, *AJ* **114**, 2366.
- CASSINELLI, J. P., MATHHIS, J. S. & SAVAGE, B. D. 1981, *Sci*, **212**, 1497.
- CHANDAR, R., BIANCHI, L., FORD, H. C., SALASNICH, B. 1999 *PASP* **111**, 794.
- CONTI, P. S., & VACCA, W. D. 1994 *ApJ* **423**, L97.
- CRABTREE, D. R. & SMECKER-HANE, T.A. 1994 *BAAS* **26**, 1494.
- DE GRIJS, R., O’CONNELL, R. W., GALLAGHER, J. S. 1999 *astro-ph/9912109*.
- DE MARCHI, G., CLAMPIN, M., GREGGIO, L., LEITHERER, C., NOTA, A. & TOSI, M. 1997, *ApJ* **479**, L27.
- DINSHAW, N., EVANS, A. S., EPPS, H., SCOVILLE, N. Z., & RIEKE, M. 1999 *ApJ* **525**, 702.
- EISENHAEUER, F., QUIRRENBACH, A., ZINNECKER, H., & GENZAL, R. 1998, *ApJ*, **498**, 278.
- ELMEGREEN, B. G. 1999, *The Evolution of Galaxies on Cosmological Timescales*, eds. J. E. Beckman & T. J. Mahoney (ASP Conference Series, San Francisco), in press.
- ELMEGREEN, B. G. & EFREMOV, Y. N. 1997 *ApJ* **480**, 235.

- FALL, S. M. & ZHANG, Q. 2000, in preparation.
- FIGER, D. F., KIM, S. S., MORRIS, M., SERABYN, E., RICH, R. M., & MCLEAN, I. S. 1999 *ApJ* **525**, 750.
- FORBES, D., BRODIE, J. P. & GRILLMAIR, C. J. 1997 *AJ* **113**, 1652.
- FORBES, D. A. & HAU, G. K. T. 2000, *MNRAS* **312**, 703.
- FRITZE-VON ALVENSLEBEN, U. 1999 *A&A* **342**, L25.
- GALLAGHER, J. S., HOMEIER, C. J., & CONSELICE, C. J. 2000a, preprint.
- GALLAGHER, S. C., HUNSBERGER, S. D., CHARLTON, J. C., ZARITSKY, D. 2000b, *Massive Star Clusters*, ed. A. Lancon & C. M. Boily (ASP conference series)
- GEORGAKAKIS, A. E., FORBES, D. A. & BRODIE, J. P. 2000, *astro-ph/0011275*.
- GILBERT, A., GRAHAM, J. R., MCLEAN, I. S., BECKLIN, E. E., LARKIN, J. WILCOX, M. K., FIGER, D. F., LEVENSON, N. A., TEPLITZ, H. I. 2000 *Massive Stellar Clusters*, ed. A. Lancon and C. M. Boily.
- GOUDFROOIJ, P., MACK, J., KISSLER-PATIG, M., MEYLAN, G., & MINNITI, D. 2000, *MNRAS*, in press.
- HARRIS, W. 1991 *ARA&A*, **29**, 543.
- HARRIS, W. 1999 private communication.
- HARRIS, W. E. & PUDRITZ, R. E. 1994 *ApJ* **429**, 177.
- HILKER, M., & M. KISSLER-PATIG 1996 *A&A* **314**, 357.
- HO, L. C. & FILIPPENKO, A. V. 1996 *ApJ* **472**, 600.
- HOLTZMAN, J. A. ET AL. (THE WFPC2 TEAM) 1996 *AJ* **112**, 416.
- HOLTZMAN, J. A. ET AL. (THE WFPC TEAM) 1992 *AJ* **103**, 691.
- HUNTER, D. A., O'CONNELL, R.W., & GALLAGHER, J. S. 1994 *AJ* **108**, 84.
- HUNTER, D. A, SHAYA, E. J., HOLTZMAN, J. A., LIGHT, R. M., O'NEIL, E. J., & LYND, R. 1995, *ApJ* **448**, 179.
- HYLAND, A. R., STRAW, S., JONES, T. J., & GATLEY, I 1992 *MNRAS* **257**, 391.
- JOG C. & SOLOMON 1992 *ApJ* **387**, 152.
- JOHNSON, K. E. & CONTI, P. 2000, *AJ* **119**, 2146.
- JOHNSON, K. E., VACCA, W. D., LEITHERER, C., CONTI, P., & LIPSEY, S. J. 1998 *AJ* **117**, 1708.
- JOHNSON, K. E., LEITHERER, C., VACCA, W. D., & CONTI, P. S. *AJ* **117**, 1708.
- KENNICUTT, R. C. & CHU. Y.-H. 1988 *AJ* **95**, 720.
- KIM, S. S., MORRIS, M., & LEE, H. M. 1999, *ApJ* **525**, 228.
- KNIERMAN, K., HUNSBERGER, S. C., GALLAGHER, S. D., CHARLTON, J. C., WHITMORE, B. C., KUNDU, A., HIBBARD, J. E., ZARITSKY, D. 2000, in preparation.
- KUMAI, Y., HASHI, Y., & FUJIMOTO, M. 1993 *ApJ* **404**, 144.
- LARSEN, S. S. & RICHTLER, T. 1999 *A&A* **345**, 59.
- LARSEN, S. S. & RICHTLER, T. 2000 *A&A* **354**, 836.
- LARSON, R. 1999 *Star Formation 1999*, ed. T. Nakamoto *astro-ph/9908189*.
- LEE, M. G., KIM, E., & GEISLER, D. 1997 *AJ* **114**, 1824.
- LEITHERER, C. & HECKMAN, T. M. 1995, *ApJS* **99**, 173.
- LEITHERER, C., VACCA, W. D., CONTI, P. S., FILIPPENKO, A. V., ROBERT, C., SARGENT, W. L. W. 1996, *AJ* **465**, 717.
- LUTZ, D. 1991 *A&A* **245**, 31.
- MALUMUTH, E. M. & HEAP, S. R. 1994, *AJ* **107**, 1054.
- MANDUSHEV, G., SPASSOVA, N. & STANEVA, A. 1991, *A&A* **252**, 94.
- MAOZ, D., BARTH, A. J., STERNBERG, A., FILIPPENKO, A. V., HO, L. C., MACCHETTO, F. D., RIX, H-W, & SCHNEIDER, D. P. 1996, *AJ* **111**, 2248.

- MASSEY, P. & HUNTER, D. 1998, *ApJ* **493**, 180.
- MENGEL, S., LEHNERT, M. D., THATTE, N., & GENZEL, R. 2000 *Massive Stellar Clusters*, ed. A. Lancon and C. M. Boily.
- MEURER, G. R. 2000 *Massive Stellar Clusters*, edited by A. Lancon and C. M. Boily (ASP).
- MEURER, G. R., FREEMAN, K. C., DOPITA, M. A. & CACCIARI, C. 1992 *AJ* **103**, 60.
- MEURER, G. R., HECKMAN, T. M., LEITHERER, C., KINNEY, A., ROBERT, C., & GARNETT, D. R. 1995 *AJ* **110**, 2665.
- MIHOS, J. C., BOTHUN, G. D., & RICHSTONE D. O. 1993 *ApJ* **418**, 82.
- MILLER, B. W., WHITMORE, B. C., SCHWEIZER, F., & FALL, S. M. 1997 *AJ* **114**, 2381.
- MIRABEL, I. F. ET AL. 1998, *A&A* **333**, L1.
- O'CONNELL, R. W., GALLAGHER, J. S., & HUNTER, D. A. 1994 *ApJ* **433**, 65.
- O'CONNELL, R. W., GALLAGHER, J. S., HUNTER, D. A., & COLLEY, W. N. 1995 *ApJ* **446**, L1.
- OSLIN, G., BERGVALL, N. RONNBACK, J. 1998, *A&A* **335**, 85
- OSLIN, G., BERGVALL, N. RONNBACK, J. 2000, *Massive Stellar Clusters*, ed. A. Lancon and C. M. Boily, (ASP Conference Series)
- PHELPS, R. L., JANES, K. A. & MONTGOMERY, K. A. 1994, *AJ* **107**, 1079.
- PORTEGIES ZWART, S. P. 2000 *astro-ph/000642*.
- RICHER, H. B., CRABTREE, D. R., FABIAN, A. C., & LIN, D. N. C. 1993 *AJ* **105**, 877.
- SALPETER, E. E. 1955 *ApJ* **121**, 161.
- SANDERS, D. B. ET AL. 1998 *ApJ* **325**, 74.
- SCALO, J. 1998 *The Stellar Initial Mass Function*, eds. G. Gilmore & D. Howell (ASP Conference Series, San Francisco), **142**, 201.
- SCHWEIZER, F. 1982 *ApJ* **252**, 455.
- SCHWEIZER, F. 1987 *Nearly Normal Galaxies*, ed. S. Faber (Springer, New York) **18**.
- SCHWEIZER, F., MILLER, B., WHITMORE, B. C., & FALL, S. M. 1996 *AJ* **112**, 1839.
- SCHWEIZER, F., & SEITZER, P. 1998 *AJ* **116**, 2206.
- SCOWEN, P. A. ET AL. 1998, *AJ* **116**, 163
- SIRIANNI, M., NOTA, A., LEITHERER, C., DE MARCHI, G. D. AND CLAMPIN, M. 2000, *AJ* **533**, 203.
- SMITH, L. & GALLAGHER, J. S. 2000 *Massive Stellar Clusters*, ed. A. Lancon and C. M. Boily.
- STIAVELLI, M., PANAGIA, N., CAROLLO, M., ROMANIELLO, M., HEYER, I., & GONZAGA, S. 1998 *ApJ* **492**, L135.
- TOOMRE, A. 1977 *The Evolution of Galaxies and Stellar Populations*, ed. B. M. Tinsley & R. B. Larson (Yale, New Haven) **401**.
- TYSON J. A. ET AL. 1998 *AJ* **116**, 102.
- VAN DEN BERGH, S. 1995 *Nature* **374**, 215.
- VAN DEN BERGH, S. 1996 *AJ* **112**, 2634.
- VAN DEN BERGH, S. 1990 *Dynamics and Interactions of Galaxies*, ed. R. Wielen (Springer, Heidelberg) **492**.
- VAN DEN BERGH S. 2000, preprint.
- VESPERINI, E. 1998 *MNRAS* **299**, 1019.
- VIGROUX, L. ET AL. 1996, *A&A* **315**, L93.
- WALBORN, N. R. & BLADES, J. C. 1997 *ApJ* **112**, 457.
- WALBORN, N. R., DRISSSEN, L., PARKER, J. W., SAHA, A., MACKENTY, J. W., & WHITE, R. L. 1999a *AJ* **118**, 1684.
- WALBORN, N. R., BARBA, R., H., BRANDNER, W., RUBIO, M., GREBEL, E., PROBST, R. 1999b *AJ* **117**, 225.

- WATSON, A. ET AL. (WFPC2 TEAM) 1996 *AJ* **112**, 534.
- WEIGELT, G. & BAIER, G. 1985, *A&A* **150**, L18
- WHITEOAK, J. B. *et al.* 1983 *MNRAS* **205**, 275.
- WHITMORE, B. C., MILLER, B. W., SCHWEIZER, F., & FALL, S. M. 1997 *AJ* **114**, 1797.
- WHITMORE, B. C., & SCHWEIZER, F. 1995 *AJ* **109**, 960.
- WHITMORE, B. C., SCHWEIZER, F., LEITHERER, C., BORNE, K., & ROBERT, C. 1993 *AJ* **106**, 1354.
- WHITMORE, B. C., ZHANG, Q., LEITHERER, C., FALL, S. M., SCHWEIZER, F., & MILLER, B. W. 1999 *AJ* **118**, 1551.
- WILSON, C. D., SCOVILLE, N., MADDEN, S. C., AND CHARMANDARIS, V. 2000 preprint
- WYSE, R. F. G., GILMORE, G., FELTZING, S., & HOUDASHELT, M. 1999 *astro-ph/9911357*
- ZEPF, S. E., & ASHMAN, K. M. 1993 *MNRAS* **264**, 611.
- ZEPF, S.E., ASHMAN, K.M., ENGLISH, J., FREEMAN, K.C., & SHARPLES, R.M. 1999 *AJ* **118**, 752.
- ZEPF, S. E., CARTER, D., SHARPLES, R. M., & ASHMAN, K. M. 1995 *ApJ* **445**, L19.
- ZHANG, Q. & FALL, S. M. 1999 *ApJ* **527**, L81.